

TOPOLOGICAL ARBITERS

MICHAEL FREEDMAN AND VYACHESLAV KRUSHKAL^{*}

ABSTRACT. This paper initiates the study of topological arbiters, a concept rooted in Poincaré-Lefschetz duality. Given an n -dimensional manifold W , a topological arbiter associates a value 0 or 1 to codimension zero submanifolds of W , subject to natural topological and duality axioms. For example, there is a unique arbiter on $\mathbb{R}P^2$, which reports the location of the essential 1-cycle. In contrast, we show that there exists an uncountable collection of topological arbiters in dimension 4. Families of arbiters, not induced by homology, are also shown to exist in higher dimensions. The technical ingredients underlying the four dimensional results are secondary obstructions to generalized link-slicing problems. For classical links in S^3 the construction relies on the existence of nilpotent embedding obstructions in dimension 4, reflected in particular by the Milnor group. In higher dimensions novel arbiters are produced using nontrivial squares in stable homotopy theory.

The concept of “topological arbiter” derives from percolation and from 4-dimensional surgery. It is not the purpose of this paper to advance either of these subjects, but rather to study the concept for its own sake. However in appendices we give both an application to percolation, and the current understanding of the relationship between arbiters and surgery. An appendix also introduces a more general notion of a multi-arbiter. Properties and applications are discussed, including a construction of non-homological multi-arbiters.

1. INTRODUCTION.

Given a closed smooth n -manifold W , consider the collection $\mathcal{M}_W$ of connected smooth codimension zero submanifolds of W .

Definition 1.1. A *topological arbiter* on W is a function $\mathcal{A}: \mathcal{M}_W \rightarrow \{0, 1\}$ satisfying axioms (1) – (3):

- (1) “ *$\mathcal{A}$ is topological*”: If $M, M' \in \mathcal{M}_W$ and M is ambiently isotopic to M' in W then $\mathcal{A}(M) = \mathcal{A}(M')$.
- (2) “*Greedy axiom*”: If $M \subset M'$ and $\mathcal{A}(M) = 1$ then $\mathcal{A}(M') = 1$.
- (3) “*Duality*”: Suppose $A, B \in \mathcal{M}_W$ are such that $W = A \cup B$, with $A \cap B = \partial A \cap \partial B$. Then $\mathcal{A}(A) + \mathcal{A}(B) = 1$.

^{*} Partially supported by the NSF.

The notion of a topological arbiter was introduced by the first author in [2] (the term *exclusive property* was used in that reference) in connection with percolation theory. The purpose of this paper is to investigate further the structure of these invariants in 4 and higher dimensions.

A prototypical example of an arbiter is the homological arbiter $\mathcal{A}_h$ on $\mathbb{R}P^2$. Specifically, given $M \subset \mathbb{R}P^2$ define $\mathcal{A}_h(M) = 1$ if M carries the non-trivial first homology class of $\mathbb{R}P^2$, i.e. if

$$H_1(M; \mathbb{Z}/2) \longrightarrow H_1(\mathbb{R}P^2; \mathbb{Z}/2) \cong \mathbb{Z}/2$$

is onto, and set $\mathcal{A}_h(M) = 0$ otherwise. The first two axioms are immediate, and (3) easily follows from Poincaré duality, see lemma 1.2 below. In fact, $\mathcal{A}_h$ is the *unique* topological arbiter on the projective plane. Indeed, suppose there is a different arbiter $\mathcal{A}$ on $\mathbb{R}P^2$, so there is $A \subset \mathbb{R}P^2$ such that $\mathcal{A}(A) = 1$ but $H_1(A; \mathbb{Z}/2) \longrightarrow H_1(\mathbb{R}P^2; \mathbb{Z}/2)$ is the trivial map. Consider the complement $B = \mathbb{R}P^2 \setminus A$, then by Poincaré-Lefschetz duality B carries the first homology of $\mathbb{R}P^2$. Since A lies in the complement of a non-trivial cycle, it is contained in a 2-cell $D^2 \subset \mathbb{R}P^2$. It follows from the axioms (2), (3) that $\mathcal{A}(D^2) = 0$, so by (2) $\mathcal{A}(A) = 0$, a contradiction proving that $\mathcal{A}_h$ is the unique arbiter on the projective plane.

It is easy to see that for any n , the n -sphere S^n does not admit a topological arbiter: consider a decomposition $S^n = A \cup B$ where both A, B are diffeomorphic to D^n . According to axiom (1), $\mathcal{A}(A), \mathcal{A}(B)$ are either both 0 or both 1, contradicting the duality axiom (3).

More generally, note that any manifold W admitting an open book decomposition does not support a topological arbiter. (However W may admit a *multi*-arbiter, see the Appendix.) Indeed, in this case the complement of the binding fibers over S^1 , where the binding is a specified codimension 2 submanifold of W . Decompose S^1 into a union of two intervals and consider the corresponding decomposition $W = A \cup B$. A and B are diffeomorphic (in fact, isotopic), so again axioms (1), (3) contradict the existence of an arbiter. Note that odd-dimensional manifolds admit open book decompositions [10, 16]. Simply-connected even-dimensional manifolds (of dimension ≥ 6) are open books, provided that the signature is zero [16]. In the general even dimensional case there is a further obstruction given by nonsingular bilinear forms [13].

Generalizing the discussion of the projective plane above, consider the “homological” arbiter on even-dimensional real and complex projective spaces:

Lemma 1.2.

(1) Let $M \subset \mathbb{R}P^{2n}$. Define $\mathcal{A}_h(M) = 0$ if $H_n(M; \mathbb{Z}/2) \longrightarrow H_n(\mathbb{R}P^{2n}; \mathbb{Z}/2) \cong \mathbb{Z}/2$ is the trivial map and $\mathcal{A}_h(M) = 1$ otherwise.

(2) Fix a field F and let $M \subset \mathbb{C}P^n$, where n is even. Define $\mathcal{A}_F(M) = 0$ if $H_n(M; F) \longrightarrow H_n(\mathbb{C}P^n; F) \cong F$ is the trivial map and $\mathcal{A}_F(M) = 1$ otherwise.

Then $\mathcal{A}_h$ is a topological arbiter on $\mathbb{R}P^{2n}$, and for any F , $\mathcal{A}_F$ is a topological arbiter on $\mathbb{C}P^n$.

Proof. The first two axioms are immediate. To prove axiom (3), let $W = \mathbb{R}P^{2n}$ or $\mathbb{C}P^n$, let the coefficients $F = \mathbb{Z}/2$ in the first case and an arbitrary field in the second case. Consider the long exact sequences

$$(1.1) \quad \begin{array}{ccccc} H_n(A) & \longrightarrow & H_n(W) & \longrightarrow & H_n(W, A) \\ \downarrow & & \downarrow & & \downarrow \\ H^n(W, B) & \longrightarrow & H^n(W) & \longrightarrow & H^n(B) \end{array}$$

where the vertical maps are isomorphisms given by Poincaré-Lefschetz duality (cf [6]). Since $H_n(W; F) \cong F$ it is clear that precisely one of the two maps $H_n(A; F) \longrightarrow H_n(W; F)$, $H_n(B; F) \longrightarrow H_n(W; F)$ is non-trivial. $\square$

There are further examples of manifolds and arbiters induced by homology. For example, generalizing lemma 1.2 one may define homological arbiters on any even-dimensional manifold W^{2n} whose middle-dimensional homology group $H_n(W; F)$ has odd rank. The arbiter $\mathcal{A}_F$ associates 1 to submanifolds representing more than half the rank of $H_n(W; F)$.

The results in this paper concern arbiters on even-dimensional balls D^{2n} , the local version of topological arbiters discussed above. *Local arbiters* (see definition 2.1 in section 2 below) associate a value 0 or 1 to codimension zero submanifolds of D^{2n} with a prescribed boundary condition: $M \cap \partial D^{2n}$ is a regular neighborhood of the standard $(n-1)$ -sphere S^{n-1} in $S^{2n-1} = \partial D^{2n}$, $\partial M \cap \partial D^{2n} = \partial D^n \times D^n$, writing $D^{2n} = D^n \times D^n$. The duality axiom satisfied by local arbiters is modeled on the Alexander duality satisfied by submanifolds $D^{2n} = A \cup B$ bounding the Hopf link $S^{n-1} \cup S^{n-1} \subset S^{2n-1}$. The reader may wish to glance ahead to absorb definition 2.1.

All arbiters discussed so far are induced by homology (with various coefficients). It follows from the universal coefficient theorem that the arbiter $\mathcal{A}_F$ on $\mathbb{C}P^n$ (and the analogous local arbiter on D^{2n}) depends only on the characteristic of the field F , so $\{\mathcal{A}_F\}$ is a countable collection. Focusing on D^4 , we construct a collection of arbiters different from the homological ones:

Theorem 1.3. *There exists an uncountable collection of local topological arbiters in dimension 4.*

The construction (discussed in more detail below and in section 2) relies on the existence of non-abelian embedding obstructions in D^4 , reflected in particular by the Milnor group. These obstructions, detecting in a certain sense the failure of the Whitney trick in 4 dimensions, have been extensively studied in the setting of link-homotopy theory [12].

Since the Whitney trick is valid in dimensions > 4 , our construction of (uncountably many) arbiters on D^4 does not have an immediate analogue in higher dimensions. In section 3 we give a rather different construction of families of local arbiters, not induced by homology, in dimensions 6 and higher. These higher-dimensional arbiters are homotopy-theoretic in nature, the following theorem summarizes our construction:

Theorem 1.4. *To each non-trivial square in the stable homotopy ring of spheres there is associated a local topological arbiter not induced by homology on D^{2n} for sufficiently large n . In particular, there exist local arbiters not induced by homology on D^{2n} for each $n > 2$.*

The main ingredient used in the construction of these arbiters in higher dimensions is the following homotopy-theoretic obstruction to a generalized link-slicing problem: for each $n > 2$ we show that there exist submanifolds $(M, S^{n-1}) \subset (D^{2n}, S^{2n-1})$, where S^{n-1} is a distinguished “attaching” sphere in the boundary of M , such that

- (1) for any coefficient ring R the attaching $(n-1)$ -sphere of M is non-trivial in $H_{n-1}(M; R)$, and
- (2) The components of the Hopf link $S^{n-1} \cup S^{n-1} \subset S^{2n-1}$ do not bound disjoint embeddings of two copies of M into D^{2n} .

Observe that if the attaching sphere S^{n-1} were trivial in $H_{n-1}(M; F)$ for some field coefficients F , then Alexander duality would immediately imply (2). The proof of (1) and (2), given in section 3, can be carried out using a “secondary” obstruction to disjointness where $n+1$ -cells are attached to S^{n-1} via the generator of π_1^S , $n > 3$. A further refinement is necessary for D^6 , see section 3.3.

To prove theorem 1.4, one defines a “partial” arbiter, setting it equal to 1 on all submanifolds of D^{2n} containing M and equal to zero on all submanifolds contained in $D^{2n} \setminus M$. The condition (2) above implies that this partial arbiter is consistently defined, and (1) shows that it is different from any homological arbiter. The proof of the theorem is completed by extending this partial arbiter to all submanifolds of D^{2n} , the details are given in section 3.

A brief outline of the rest of the paper is as follows. The proof of theorem 1.3 begins in section 2 with the introduction of a dyadic collection of “model” submanifolds of D^4 (i.e. a special collection of submanifolds, parametrized by vertices of a trivalent tree.) A generalization of the Milnor group is used to find obstructions to pairwise embeddings of these model submanifolds. The collection $\mathcal{M}_{D^4}$ of all submanifolds of D^4 has a partial ordering induced by inclusion. The non-embedding results mentioned above allow one to define an uncountable collection of arbiters on those elements of $\mathcal{M}_{D^4}$ which are comparable to model submanifolds. One then extends these arbiters to $\mathcal{M}_{D^4}$, concluding the proof of theorem 1.3, see section 2.7.

Section 2.9 points out that the embeddings of the submanifolds $M \hookrightarrow W$ in definition 1.1 of arbiters play a very important role. The result of [8] shows that if one considers topological arbiters as invariants of manifolds, disregarding their embedding information, then in the context of D^4 whenever the attaching circle of $M \subset D^4$ is trivial in $H_1(M; \mathbb{Z})$ the value of any arbiter on M has to be 1. The proof of theorem 1.3 is valid only when the embedding data is taken into account. Indeed the arbiters constructed in its proof assume value 1 on manifolds where the attaching circle does not vanish in homology with any coefficients.

Section 2.10 adapts the local construction in the proof of theorem 1.3 to the setting of $\mathbb{C}P^2$. It seems likely that this construction should yield an uncountable collection of arbiters on $\mathbb{C}P^2$, however proving this would require a substantial extension of the techniques of Milnor's theory of link homotopy.

In appendix A we note that the existence of a local arbiter on D^4 satisfying an additional ‘‘Bing doubling’’ axiom would give an obstruction to topological 4-dimensional surgery for the non-abelian free groups, using the A-B slice reformulation [3, 4] of the surgery conjecture.

Appendix B introduces two generalizations of topological arbiters, *multi-arbiters*. We discuss their properties and applications, in particular a construction of non-homological multi-arbiters, and an application to percolation theory.

2. LOCAL ARBITERS.

The purpose of this section is to prove theorem 1.3. We start by defining the notion of a local topological arbiter on D^4 (more generally, the definition makes sense for D^{2n} for any even dimension $2n$.) Consider $\mathcal{M} = \mathcal{M}_{D^4} = \{(M, \gamma) \mid M \text{ is a codimension zero, smooth, compact submanifold of } D^4, \gamma \subset \partial M, \text{ and } M \cap \partial D^4 \text{ is a tubular neighborhood of an unknotted circle } \gamma \subset S^3\}$. (For $n > 2$, $M \cap \partial D^{2n}$ is a tubular neighborhood of a standard S^{n-1} , isotopic in $\partial D^{2n} = \partial(D^n \times D^n)$ to $\partial D^n \times D^n$.) Consider a local version of definition 1.1:

Definition 2.1. A *local topological arbiter* is a function $\mathcal{A}: \mathcal{M} \rightarrow \{0, 1\}$ satisfying axioms (1) – (3):

- (1) If $(M, \gamma), (M', \gamma') \in \mathcal{M}$ and (M, γ) is ambiently isotopic (not necessarily relative to the identity on the boundary) to (M', γ') in (D^4, S^3) then $\mathcal{A}(M, \gamma) = \mathcal{A}(M', \gamma')$.
- (2) If $(M, \gamma) \subset (M', \gamma')$ and $\mathcal{A}(M, \gamma) = 1$ then $\mathcal{A}(M', \gamma') = 1$.
- (3) Suppose $(A, \alpha), (B, \beta) \in \mathcal{M}_{D^4}$ are such that $D^4 = A \cup B$, with $A \cap B = \partial A \cap \partial B$, and so that the attaching curves α, β of A, B form the Hopf link in ∂D^4 . (In other words, $D^4 = A \cup B$ is a decomposition of D^4 extending the standard genus 1 Heegaard decomposition of the 3–sphere.) Then $\mathcal{A}(A, \alpha) + \mathcal{A}(B, \beta) = 1$. (For $n > 2$ the ‘‘Hopf link’’ above is replaced with $\partial D^n \times 0 \sqcup 0 \times \partial D^n \subset S^{2n-1}$.)

We summarize local versions of the properties discussed in the introduction. In dimension 4 (more generally, in any even dimension), given a field F there is a “homological” local arbiter $\mathcal{A}_F$. This arbiter (by the universal coefficient theorem depending only on the characteristic of F) is defined by $\mathcal{A}_F(M, \gamma) = 1$ if and only if $\gamma = 0 \in H_1(M; F)$. The axioms (1), (2) are satisfied automatically, (3) is a consequence of Alexander duality.

In dimension 2 there is a unique local arbiter $\mathcal{A}$ (equal to $\mathcal{A}_F$ for any field F .) The difference between arbiters in dimensions 2 and higher is explained by the divergence between homology and homotopy: for $(A, \alpha) \subset (D^2, S^1)$, $\alpha = 0 \in H_0(A)$ iff $\alpha = 0 \in \pi_0(A)$. If this is the case, the complement B of A in D^2 may be pushed off an arc connecting α in A into a collar on its attaching curve β , therefore the value of any arbiter on B is zero. In higher dimensions, being trivial in homology is a weaker condition, and we show that there exist arbiters whose value on B may be 1 even though β is homologically non-trivial with any coefficients.

2.2. Model decompositions. The proof of theorem 1.3 starts with the construction of a special collection of submanifolds of D^4 , which are shown below to satisfy pairwise non-embedding properties. These codimension zero submanifolds form complementary pairs A_I, B_I parametrized by a dyadic multiindex I discussed in more detail below. For each index I , the pair is a decomposition of the 4-ball in the sense of definition 2.1 (3) above: $D^4 = A_I \cup B_I$, and the distinguished curves α_I, β_I form the Hopf link in ∂D^4 . To start the construction, consider the genus one surface S with a single boundary component, and set $B_1 = S \times D^2$. Denote $\beta_1 = \partial S \times 0$. To specify an embedding $(B_1, \beta_1) \subset (D^4, S^3)$, embed the surface S into S^3 , push it into the 4-ball and take a regular neighborhood.




FIGURE 1.

Proposition 2.3. *The complement $A_1 = D^4 \setminus B_1$ is obtained from the collar $S^1 \times D^2 \times I$ by attaching a pair of zero-framed 2-handles to the Bing double of the core of the solid torus $S^1 \times D^2 \times 1$, figures 1, 2.*

The distinguished curve $\alpha_1 \subset \partial A_1$ is formed by $S^1 \times 0 \times 0$. Figure 1 is a schematic “spine” picture of the manifolds A_1, B_1 , a precise description in terms of Kirby diagrams (drawn in a solid torus) is given in figure 2. Note that a distinguished pair




FIGURE 2.

of curves γ_1, γ_2 , forming a symplectic basis in the surface S , is determined as the meridians (linking circles) to the cores of the 2–handles H_1, H_2 in D^4 . In other words, γ_1, γ_2 are fibers of the circle normal bundles over the cores of H_1, H_2 in D^4 . The proof of proposition 2.3 is an exercise in Kirby calculus, see for example [5].

The manifolds A_1, B_1 are the basic building blocks in the inductive construction of model decompositions. Consider the 2-handle H_1 (one of the 2-handles in the definition of A_1 , figure 1) in place of the original 4-ball. The pair of curves (γ_1 , the attaching circle δ_1 of H_1) forms the Hopf link in the boundary of H_1 . In H_1 consider the standard genus one surface bounded by δ_1 . As discussed above, its complement is given by two zero-framed 2-handles attached to the Bing double of γ_1 . Assembling this data, consider the new decomposition $D^4 = A_0 \cup B_0$. The manifold A_0 and its Kirby diagram are shown in figure 7 below, and both A_0, B_0 are included in figure 3.

We are now in a position to define the more general collection of special elements A_I, B_I of $\mathcal{M}$ (“model decompositions”, $D^4 = A_I \cup B_I$), figure 3. These decompositions correspond to vertices of a trivalent tree, and each A_I, B_I is labeled by a dyadic multiindex I . Each arrow in figure 3 represents a an inclusion. These model decompositions are defined inductively; the transition from A_I to $A_{I,0}$ replaces one of the two 2-handles of A_I , introduced during the previous step, by A_0 . $A_{I,1}$ is obtained by replacing one of these two “new” 2-handles of A_I by A_1 (in other words, by Bing doubling this 2-handle.) Then B_I is defined as the complement $D^4 \setminus A_I$. Note that this is a subset of the collection of decompositions of D^4 introduced in the context of the A-B slice problem in [5]. Figure 3 shows only the “spines” of the submanifolds A_I, B_I ; we refer the reader to [5] for a precise description of these 4-manifolds in terms of Kirby diagrams.

Consider the *lexicographic order* on the set of dyadic multiindices: given $I = (i_1, \dots, i_m)$, $J = (j_1, \dots, j_n)$, set $I < J$ if $i_1 = j_1, \dots, i_k = j_k$ and $i_{k+1} = 0, j_{k+1} = 1$ for some k . Also set $I < J$ if I is an initial segment of J : $J = (I, j_{k+1}, \dots)$

Consider the trivalent tree T encoding the submanifolds A_I . The root of T corresponds to $A = 2$ -handle. Let r be a “geodesic” ray in T originating at the root; r is viewed as an infinite dyadic multiindex. For a technical reason which will be




FIGURE 3. The collection of model decompositions A_I, B_I . The arrows represent inclusions.

apparent in the proof below, it is convenient to consider all rays r passing through

A_1 (i.e. whose first coefficient is 1), and the lexicographic order is extended to the set of such rays.

Definition 2.4. Let $\mathcal{M}_0 \subset \mathcal{M}$ denote the collection of submanifolds $\{A_I, B_I\}$ defined above. For each r , a ray as above, define a *partial arbiter* $\mathcal{A}_r: \mathcal{M}_0 \rightarrow \{0, 1\}$ by setting

$$\mathcal{A}_r(A_I) = 1 \text{ if } I < r, \text{ and } \mathcal{A}_r(A_I) = 0 \text{ if } I > r.$$

Then the duality axiom (3) in definition 2.1 forces one to define

$$\mathcal{A}_r(B_I) = 0 \text{ if } I < r, \text{ and } \mathcal{A}_r(B_I) = 1 \text{ if } I > r.$$

Lemma 2.5. For each r , $\mathcal{A}_r: \mathcal{M}_0 \rightarrow \{0, 1\}$ satisfies the greedy axiom (2) in definition 2.1. In other words, given $(M, \gamma), (M', \gamma') \in \mathcal{M}_0$ with $(M, \gamma) \subset (M', \gamma')$, if $\mathcal{A}_r(M, \gamma) = 1$ then $\mathcal{A}_r(M', \gamma') = 1$.

The proof follows from the following proposition. Recall from the beginning of section 2 that each element $(M, \gamma) \in \mathcal{M}$ comes equipped with an embedding $(M, \gamma) \subset (D^4, S^3)$.

Proposition 2.6.

- (1) If $I < J$ then $(A_I, \alpha_I) \not\subset (A_J, \alpha_J)$.
- (2) For any I, J , $(A_I, \alpha_I) \not\subset (B_J, \beta_J)$.
- (3) For any multiindices I, J whose first entries equal 1, there does not exist an embedding $(B_I, \beta_I) \subset (A_J, \alpha_J)$.

We postpone the proof of proposition 2.6 to complete the proofs of lemma 2.5 and theorem 1.3. One needs to check that for any two elements $(M_1, \gamma_1), (M_2, \gamma_2) \in \mathcal{M}_0$ if $\mathcal{A}_r(M_1) = 1$ and $\mathcal{A}_r(M_2) = 0$ then $(M_1, \gamma_1) \not\subset (M_2, \gamma_2)$. Consider the list of all pairs M_1, M_2 where $\mathcal{A}_r(M_1) = 1$ and $\mathcal{A}_r(M_2) = 0$:

- (a) $M_1 = A_I$ where $I < r$, $M_2 = A_J$ where $J > r$,
- (b) $M_1 = A_I$ where $I < r$, $M_2 = B_J$ where $J < r$,
- (c) $M_1 = B_I$ where $I > r$, $M_2 = A_J$ where $J > r$,
- (d) $M_1 = B_I$ where $I > r$, $M_2 = B_J$ where $J < r$.

The cases (a), (b), (c) follow from the statements (1), (2), (3) of proposition 2.6 respectively. To eliminate (d), suppose $B_I \subset B_J$ where $J < I$. Considering their complements, there is an inclusion $A_J \subset A_I$, a contradiction with proposition 2.6 (1). This completes the proof of lemma 2.5. $\square$




FIGURE 4. The vertices of the graph encode codimension zero submanifolds of D^4 (elements of $\mathcal{M}$), and the edges correspond to inclusions. This figure illustrates the case $I < r$, $J > r$: then $\mathcal{A}_r(A_I) = \mathcal{A}_r(B_J) = 1$, and $\mathcal{A}_r(B_I) = \mathcal{A}_r(A_J) = 0$. These requirements force $\mathcal{A}_r(M) = 1$ for any M in the “cones” over A_I, B_J and $\mathcal{A}_r(M) = 0$ for any M in the “cones” under B_I, A_J . The upward-facing cones are disjoint from the down-facing cones by lemma 2.5.

2.7. Proof of theorem 1.3. Given a ray r , by lemma 2.5 the partial arbiter $\mathcal{A}_r$ is consistently defined on the collection $\mathcal{M}_0$ of model decompositions. Given $M_1, M_2 \in \mathcal{M}_0$ with $\mathcal{A}_r(M_1) = 0$, $\mathcal{A}_r(M_2) = 1$, axiom (2) in definition 2.1 forces $\mathcal{A}_r(M) = 0$ for any M isotopic to a submanifold of M_1 and $\mathcal{A}_r(M) = 1$ for any M isotopic to M' where $M_2 \subset M'$. Graphically, $\mathcal{A}_r = 0$ in the “cone” below M_0 and $\mathcal{A}_r = 1$ in the “cone” above M_1 , figure 4. The upward-facing cones are disjoint from the down-facing cones by lemma 2.5. This defines a consistent extension of $\mathcal{A}_r$ to the union of all such cones. Duality axiom (3) in definition 2.1 for this extension follows from the fact that the upward-facing cone over, say, A_I consists precisely of the complements in D^4 of the elements of the down-facing cone under B_I . To extend $\mathcal{A}_r$ to $\mathcal{M}$, consider any local arbiter on D^4 , for example a homological arbiter $\mathcal{A}_F$ for some fixed field F . Set $\mathcal{A}_r(M) = \mathcal{A}_F(M)$ for any M in the complement of the union of the “cones” discussed above. It is straightforward to check that $\mathcal{A}_r$ satisfies the axioms of a local arbiter.

Note that if $r \neq r'$ then $\mathcal{A}_r \neq \mathcal{A}_{r'}$: assume $r < r'$, then their difference is detected by a codimension zero submanifold A_I where I is a multiindex with $r < I < r'$. This completes the proof of theorem 1.3, assuming proposition 2.6. $\square$

Remarks.

1. Because we use a definite procedure to extend $\mathcal{A}_r$ beyond the “cones” on which it is determined by r , the proof of theorem 1.3 does not depend on the axiom of choice. Nor does any other assertion in this paper. To be complete, our argument actually has shown that the cardinality of local arbiters on D^4 is precisely the continuum.

2. The argument extending partial arbiters $\mathcal{A}_r$ to arbiters defined on all codimension zero submanifolds of D^4 in the proof of theorem 1.3 may also be used to define “hybrids” of homological arbiters of different characteristics. For example, suppose (M, γ) is an element of $\mathcal{M}$ such that $\gamma = 0 \in H_1(M; \mathbb{Z}/p)$ but $\gamma \neq 0 \in H_1(M; \mathbb{Z}/q)$ for some primes p and q . By Alexander duality with $\mathbb{Z}/p$ coefficients, two copies of M cannot be embedded disjointly into D^4 so that the distinguished curves form the Hopf link on the boundary. Then a partial arbiter $\mathcal{A}$ may be defined by setting it equal to $\mathcal{A}_{\mathbb{Z}/p}$ in the cone over M and also in the cone under $D^4 \setminus M$. Extend it by $\mathcal{A}_{\mathbb{Z}/q}$ to the rest of $\mathcal{M}$. The arbiters constructed in the proof of theorem 1.3 assume value 1 on elements $(M, \gamma) \in \mathcal{M}$ such that $\gamma \neq 0 \in H_1(M; R)$ for **any** coefficients R , therefore they are different from any such hybrid of homological arbiters.

Proof of part (1) of proposition 2.6. Before giving the general (inductive) proof consider several special cases which clarify the inductive step. (Refer to figure 3 and the beginning of section 2.2 to recall the definition of the submanifolds A_I, B_I .)

(i) (A_0, α_0) does not embed in (A_1, α_1) .

The proof is immediate since $\alpha_0 = 0 \in H_1(A_0)$ and $\alpha_1 \neq 0 \in H_1(A_1)$.

(ii) $(A_{10}, \alpha_{10}) \not\subset (A_{11}, \alpha_{11})$.

Suppose such an embedding exists. Then consider the complements: $(B_{11}, \beta_{11}) \subset (B_{10}, \beta_{10})$. The submanifold B_{11} is (a thickening of) a 2-stage *grope*, see [5] (the term used in that reference is a *half-grope*), a geometric counterpart of the commutators in the lower central series π_n of a group π . This series is defined inductively by $\pi_n = [\pi, \pi_{n-1}]$, $\pi_1 = \pi$. The attaching curve β_{11} of the 2-stage grope B_{11} is in the third term $(\pi_1(B_{11}))_3$. This is a contradiction since β_{11} maps to β_{10} , however

$$\beta_{11} = 1 \in \pi_1(B_{11})/(\pi_1(B_{11}))_3 \text{ but } \beta_{10} \neq 1 \in \pi_1(B_{10})/(\pi_1(B_{10}))_3.$$

(iii) $(A_{00}, \alpha_{00}) \not\subset (A_{01}, \alpha_{01})$.

The argument in this example will serve as an ingredient in the proof in the general case further below. Suppose the embedding i exists as in figure 5. Considering the complements one gets $\bar{i}: (B_{01}, \beta_{01}) \rightarrow (B_{00}, \beta_{00})$, figure 6. The argument will involve two steps, the first one using the embedding i and second step using $\bar{i}$. The connection between the two steps is provided by the Alexander duality.

Step 1. Denote by Σ', Σ'' the bottom surface stages of A_{00}, A_{01} . Since α_{00} maps to α_{01} , i “restricts” to a degree 1 map $(\Sigma', \alpha_{00}) \rightarrow (\Sigma'', \alpha_{01})$. (More precisely,




FIGURE 5.




FIGURE 6.

consider the restriction $i|_{\Sigma'}$. A_{01} is homotopy equivalent to the wedge of Σ'' and three 2-spheres. The composition of $i|_{\Sigma'}$ with the retraction onto Σ'' has degree 1.)

Consider the classes in $H_2(B_{00})$, $H_2(B_{01})$ represented by the tori T' , T'' and spheres S' , S'' , figure 6. (These are closed surfaces since they are capped off by the null-homotopies for the Bing-double curves.) The punctured tori $\Sigma' \subset A_{00}$, $\Sigma'' \subset A_{01}$ have preferred symplectic bases of curves: γ_1, γ_2 on Σ' , and δ_1, δ_2 on Σ'' . The curves γ_1, γ_2 are Alexander dual respectively to T', S' , and similarly δ_1, δ_2 are dual to T'', S'' . Consider the matrix $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ representing the induced map $i_*: H_1(\Sigma') \rightarrow H_1(\Sigma'')$ in these bases. Since the map $(\Sigma', \partial) \rightarrow (\Sigma'', \partial)$ has degree 1, M is an element of $SL_2(\mathbb{Z})$.

Step 2. The matrix for the map $\bar{i}_*: H_2(B_{01}) \rightarrow H_2(B_{00})$, restricted to the 2-dimensional subspaces spanned by $[S''], [T'']$ and $[S'], [T']$, is the transpose of M , by Alexander duality. Since $[S'']$ is a spherical class, the component of $\bar{i}_*([S''])$ along $[T']$ is trivial, so $b = 0$ and M must be of the form $\begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix}$. Therefore the class $[T'']$ maps to $[T']$ plus a multiple c of $[S']$, but in particular one gets a degree 1 map

$T'' \longrightarrow T'$. This leads to a contradiction when one considers the elements represented by T' , T'' in the *Dwyer's filtration* (see [1]) of the corresponding spaces. A geometric description of the n th term of the Dwyer's filtration $\phi_n(X) \subset H_2(X; \mathbb{Z})$ of a space X is the subgroup consisting of those elements in $H_2(X; \mathbb{Z})$ which may be represented by maps of closed gropes of class n into X . Since $[T''] \in \phi_3(B_{01})$ but $[T'] \notin \phi_3(B_{00})$, $[T'']$ cannot map to $[T']$, a contradiction.

Proof of part (1) of proposition 2.6 in the general case. Suppose an embedding $i: (A_I, \alpha_I) \longrightarrow (A_J, \alpha_J)$ exists; let $\bar{i}: (B_J, \beta_J) \longrightarrow (B_I, \beta_I)$ be the corresponding inclusion of the complements. Since $I < J$, $I = (i_1, \dots, i_n, 0, \dots)$ and $J = (i_1, \dots, i_n, 1, \dots)$ for some n .

First consider the case $i_1 = \dots = i_n = 0$, the proof for arbitrary $i_1, \dots, i_n$ will be analogous. Since all i_j are assumed to be zero, $j = 1 \dots, n$, each of the first n steps in the construction of A_I, A_J replaces an existing 2-handle with (a thickening of) the genus 1 surface, and then attaches a pair of 2-handles to the Bing double of an essential curve on this surface. Denote the successive genus 1 surface stages of A_I by $\Sigma'_1, \dots, \Sigma'_n$, and similarly denote the genus 1 surface stages of A_J by $\Sigma''_1, \dots, \Sigma''_n$. The construction on the B -side is dual (in the sense of proposition 2.3): each of the first n steps attaches a pair of 2-handles to a Bing double, and then replaces one of these 2-handles with a punctured torus. Denote the successive tori in B_I by $T'_1, \dots, T'_n$ and the corresponding tori in B_J by $T''_1, \dots, T''_n$. (In the special case $I = (00), J = (01)$ considered in (iii) above, n was equal to 1.)

The base of the induction. The embedding i induces a degree 1 map $(\Sigma'_1, \alpha_I) \longrightarrow (\Sigma''_1, \alpha_J)$.

The inductive step. The inductive hypothesis is that the embedding i induces a degree 1 map $\Sigma'_k \longrightarrow \Sigma''_k$ for some $1 \leq k < n$. The argument in the special case (iii) above proves that then the complementary embedding $\bar{i}$ induces a degree 1 map $T'_k \longrightarrow T''_k$. Since the construction of the first n stages of the submanifold A_I is precisely dual to the construction of B_I (and similarly for A_J, B_J , compare figures 5, 6), exactly the same argument now propagates the conclusion to the A -side: if $\bar{i}$ induces a degree 1 map $T''_k \longrightarrow T'_k$, then the embedding i induces a degree 1 map $\Sigma'_{k+1} \longrightarrow \Sigma''_k$. This concludes the proof of the inductive step.

A contradiction is reached after $n - 1$ applications of the inductive step: the outcome is a degree 1 map $\Sigma'_n \longrightarrow \Sigma''_n$; run the argument one more time to get a degree 1 map $T''_n \longrightarrow T'_n$. However since the $n + 1$ -st coefficient of J is 1, it follows from the construction of B_J that $[T''_n] \in \phi_3(B_J)$. On the other hand the $n + 1$ -st coefficient of I is 0, so $[T'_n] \notin \phi_3(B_I)$. This contradicts the fact that $\bar{i}$ induces a degree 1 map $T''_n \longrightarrow T'_n$, concluding the proof when $i_1 = \dots = i_n = 0$.

A slight modification of the argument is necessary in the general case when $i_1 = 0$ but $(i_1, \dots, i_n)$ contains subsequences with one or more consecutive 1's. Then the

corresponding tori on the B -side are replaced by gropes of height h equal to $(1 + \text{the number of consecutive } 1\text{'s in a subsequence})$, all of whose surface stages have genus 1. (Correspondingly, the 2-handles on the A -side are attached to higher iterated Bing doubles.) The inductive step discussed above applies without any changes on the A -side.

On the B -side a variation is necessary: Let f be a map $f: G \rightarrow G$ where G is a half-grope of height n with all surface stages of genus 1. The second homology class of the bottom stage surface of G reads off the relation $[\gamma_1, [\gamma_2, \dots, [\gamma_h, \gamma_{h+1}] \dots]]$, where γ_i are the curves generating $\pi_1(G)$, corresponding to the “tips” of the grope. Note that γ_h, γ_{h+1} generate π_1 of the top stage surface of G . Suppose f_* carries the class of the bottom stage surface of G in $H_2(G)$ to itself. Then the map induced by f on the 2-dimensional subspace of $H_1(G)$ spanned by γ_h, γ_{h+1} is in $SL(2, \mathbb{Z})$. The rest of the argument in the inductive step goes through.

Finally, if $i_1 = 1$ then the first step of the induction is applied on the B -side. This concludes the proof of 2.6 (1).

The proof of part (2) of proposition 2.6 involves a generalization of the Milnor group from the context of link homotopy (a link in S^3 bounding disjoint maps of disks in D^4) to a class of more general submanifolds of the 4-ball. We outline this construction below and refer the reader to [9] for a detailed exposition.

Suppose $(A_I, \alpha_I) \subset (B_J, \beta_J)$. Then A_I embeds in the complement of A_J , where the attaching curves α_I, α_J form the Hopf link in $S^3 = \partial D^4$. As a warmup, consider the special case where both I, J are dyadic multiindices with no zeros, for example A_1 and A_{11} are shown in figure 3. For such multiindices I, J , the handlebodies A_I, A_J consist of 2-handles attached to iterated Bing doubles of their respective attaching curves α_I, α_J which form the Hopf link in ∂D^4 . It is straightforward to prove that any link L obtained by iterated Bing doubling of the Hopf link is a homotopically essential link [12], by computing that it has a non-trivial Milnor’s $\bar{\mu}$ -invariant with non-repeating coefficients. Therefore the components of L do not even bound disjoint maps of disks into D^4 , contradicting the assumption $(A_I, \alpha_I) \subset (B_J, \beta_J)$.

Now consider a more interesting example A_0 (figure 7), illustrating the argument in the general case. Consider a pair of pants P with boundary components $\gamma, \alpha_1, \alpha_2$, and let C denote $P \times D^2 \cup$ four zero-framed 2-handles $h_1, \dots, h_4$ attached to the Bing doubles of the curves α_1, α_2 , figure 8.

Observe that there is a map $C \rightarrow A_0$ taking γ to α_0 , such that the 2-handle h_1 is disjoint from h_2 , and similarly h_3 is disjoint from h_4 . On the level of spines, the proof amounts to cutting the punctured torus into a pair of pants; the separation of parallel copies of the 2-handles results in $h_1 - h_3$ and $h_2 - h_4$ double points.

Therefore to prove that the components of the Hopf link do not bound disjoint embeddings of two copies of A_0 , it suffices to show that its components do not bound




FIGURE 7. The submanifold A_0 and its Kirby diagram. (The handle diagram is drawn in the solid torus which is the complement of the dashed circle in the 3-sphere.)




FIGURE 8. The handlebody C (a Bing cell of height 1): a schematic picture and a Kirby diagram

disjoint maps of two copies C', C'' of C , subject to the disjointness condition on the 2-handles discussed above. Suppose the opposite is true. Consider meridians m'_i to the 2-handles $\{h'_i\}$ of C' and meridians m''_i to the 2-handles $\{h''_i\}$ of C'' in D^4 . (By a meridian we mean a based loop in the complement representing the same homology class as a fiber of the normal circle bundle over a 2-handle.) Consider the quotient $\overline{M}\pi$ of $\pi := \pi_1(D^4 \setminus (C' \cup C''))$ by the relations $[(m'_i)^x, (m'_j)^y]$ and $[(m''_i)^x, (m''_j)^y]$, where $\{i, j\} \neq \{1, 2\}, \{3, 4\}$ and the conjugating elements x, y range over all elements of π . These relations correspond to the Clifford tori linking the double points between the various 2-handles in D^4 , and the fact that the pairs $\{1, 2\}, \{3, 4\}$ are excluded reflects the fact that these handles are required to be disjoint. (This is a generalization of the Milnor group $M\pi$ (see [12]) which in this context is defined as the quotient of π by the relations $[(m'_i)^x, (m'_i)^y], [(m''_i)^x, (m''_i)^y], i = 1, \dots, 4$, which correspond to *self*-intersections of the handles. These relations are included in the defining relations of $\overline{M}\pi$.)

The classical Milnor group $M\pi$ is nilpotent, and since $\overline{M}\pi$ is a quotient of $M\pi$, it is nilpotent as well. It follows that $\overline{M}\pi$ is generated by the chosen meridians, and its relations may be read off from the generators of $H_2(D^4 \setminus (C' \cup C''); \mathbb{Z})$, which is Alexander dual to the relative group $H_1(C' \cup C'', \partial(C' \cup C'') \cap \partial D^4)$. There are two types of generators of this second homology group: the Clifford tori that give rise to the commutation relations used in the definition of $\overline{M}\pi$, and the generators dual to the relative first homology group of the pairs of pants defining C', C'' . The latter relations are of the form $[m'_1, m'_2] = [m'_3, m'_4], [m''_1, m''_2] = [m''_3, m''_4]$.

Let m', m'' denote the respective meridians to the components of the Hopf link in ∂D^4 , formed by the attaching curves γ', γ'' . The relation $[m', m''] = 1$ holds in $\pi_1(S^3 \setminus (\gamma' \cup \gamma''))$, therefore it also holds in $\pi_1(D^4 \setminus (C' \cup C''))$. On the other hand, in terms of the generators of $\overline{M}\pi$, $m' = [m'_1, m'_2] = [m'_3, m'_4], m'' = [m''_1, m''_2] = [m''_3, m''_4]$. One then shows using the Magnus expansion that, based on the relations in $\overline{M}\pi$ discussed above, the commutator $[m', m'']$ is in fact non-trivial in $\overline{M}\pi$: phrased differently, there are “not enough” relations in $\pi_1(D^4 \setminus (C' \cup C''))$ to imply the relation $[m', m''] = 1$ which holds in $\pi_1(S^3 \setminus (\gamma' \cup \gamma''))$. This contradiction shows that the components of the Hopf link do not bound in D^4 disjoint copies of the manifold A_0 .

The manifolds A_0, A_1 considered above are the building blocks in the definition of model decompositions $D^4 = A_I \cup B_I$, see section 2.2. There is a similar inductive construction of *Bing cells* based on the manifold C above, where the inductive step replaces a 2-handle with a copy of C . *Moreover, the 2-handles in the more general definition below may be attached to iterated Bing doubles*, since such links arise in the model submanifolds A_I (see for example A_{01} in figure 3.)

Definition 2.8. A *model Bing cell of height 1* is a smooth 4-manifold C with boundary and with a specified curve $\gamma \subset \partial C$, defined as follows. Consider a pair of pants P with boundary components $\gamma, \alpha_1, \alpha_2$, and let L_i be a (possibly iterated) Bing double of the core of the solid torus $\alpha_i \times D^2$. Then C is obtained from $P \times D^2$ by attaching zero-framed 2-handles along the components of $L_1 \cup L_2$.

A *model Bing cell C of height h* is obtained from a model Bing cell of height $h - 1$ by replacing some of its 2-handles with Bing cells of height one.

Bing cells in a 4-manifold M are defined as maps of model Bing cells into M , subject to certain crucial disjointness requirements. For each link L_i in the inductive definition above, surfaces and more generally Bing cells attached to different components of L_i are required to be disjoint in M . Generalizing the arguments above, one shows (see [9]) that

- (1) If the components of a link $L \subset S^3 = \partial D^4$ bound disjoint Bing cells in D^4 then L is homotopically trivial, and
- (2) For each index I the attaching curve α_I bounds a Bing cell in A_I .

Therefore the components of the Hopf link do not bound disjoint embeddings of A_I, A_J , i.e. $(A_I, \alpha_I) \not\subset (B_J, \beta_J)$. This concludes the proof of proposition 2.6 (2).

The proof of part (3) of proposition 2.6 is immediate since $\beta_I = 0 \in H_1(B_I)$ and $\alpha_J \neq 0 \in H_1(A_J)$. This completes the proof of proposition 2.6 and of theorem 1.3. $\square$

2.9. Arbiters and embeddings. In this section we note the important role played by the embeddings in definition 2.1 of arbiters on D^4 . According to [8] for any given g there exist decompositions (in the sense of axiom (3) in definition 2.1) $D^4 = A_g \cup B_g$, where the attaching curves α, β form the Hopf link in ∂D^4 , such that the conditions (i), (ii) are satisfied:

- (i) Let S denote the genus g surface with one boundary component, $\gamma = \partial S$. Denote by S_0 its untwisted 4-dimensional thickening, $S_0 = S \times D^2$, and set $\gamma_0 = \gamma \times \{0\}$. Then there exists a proper embedding $(A_g, \alpha) \subset (S_0, \gamma_0)$.
- (ii) B_g embeds in a collar on its attaching curve. More precisely, there exists a proper embedding $(B_g, \beta) \subset (S^1 \times D^2 \times [0, 1], S^1 \times \{0\} \times \{0\})$

Suppose $\mathcal{A}$ is an invariant as in definition 2.1, where the embedding in the axiom (2) in 2.1 is assumed to be an arbitrary embedding of abstract pairs $(M, \gamma), (M', \gamma')$, rather than just the inclusion of submanifolds of D^4 . Since the value of $\mathcal{A}$ on the collar is necessarily 0, (ii) implies that for each g , $\mathcal{A}(B_g) = 0$, therefore $\mathcal{A}(A_g) = 1$. (Note that the embeddings in (i), (ii) are indeed not isotopic to the original embeddings into D^4 .) Now suppose (M, γ) is a submanifold of (D^4, S^3) such that the attaching curve γ of the 4-manifold M is trivial in $H_1(M; \mathbb{Z})$. Then γ bounds an embedded surface S in M with the trivial normal bundle. It follows that $(A_g, \alpha) \subset (M, \gamma)$, where g is the genus of S . Again by axiom (2) which allows arbitrary embeddings, $\mathcal{A}(M, \gamma) = 1$. That is, a function $\mathcal{A}$ which is insensitive to the embedding of submanifolds into D^4 is determined by homology, at least on those submanifolds whose attaching curve is trivial in the first homology with integer coefficients, and on their complements in D^4 . The proof of theorem 1.3 is valid only when the embeddings are inclusions of submanifolds of D^4 , indeed the arbiters constructed in its proof assume value 1 on manifolds where the attaching circle does not vanish in homology with any coefficients.

2.10. Submanifolds of $\mathbb{C}P^2$. This section outlines a construction of submanifolds of $\mathbb{C}P^2$ which may be viewed as an “absolute” analogue of the relative model decompositions of D^4 defined in section 2.2.

Consider the usual handle structure on $\mathbb{C}P^2$ with one handle of each index 0, 2, 4: $\mathbb{C}P^2 = 0H \cup 2H \cup 4H$. Consider decompositions $\mathbb{C}P^2 = \hat{A} \cup \hat{B}$, where $\hat{A} \cap \hat{B} = \partial \hat{A} \cap \partial \hat{B}$. The decompositions are constructed of the form $\hat{A} = 0H \cup A$, $\hat{B} = 4H \cup B$, and $2H = A \cup B$ is a decomposition of the 4-ball $2H = D^2 \times D^2$ in the sense

discussed in section 2: $A \cup B$ extends the standard genus 1 decomposition of the 3-sphere $D^2 \times S^1 \cup S^1 \times D^2$.

Applying this construction to model decompositions $D^4 = A_I \cup B_I$ of the 4-ball in section 2.2 (see figure 3), one gets a dyadic collection of decompositions $\mathbb{C}P^2 = \widehat{A}_I \cup \widehat{B}_I$. More concretely, $\widehat{A}_I$ is obtained by attaching a 1-framed 2-handle to A_I along the attaching curve α_I , and similarly its complement $\widehat{B}_I$ is obtained by attaching a 1-framed 2-handle to B_I along the attaching curve β_I .

It seems reasonable to conjecture that there are non-embedding results analogous to proposition 2.6, which should enable one to construct an uncountable collection of topological arbiters on $\mathbb{C}P^2$. However proving this would require an extension of the Milnor group techniques which is outside the scope of this paper. This question remains open.

3. SQUARES IN STABLE HOMOTOPY AND ARBITERS IN HIGHER DIMENSIONS

Recall that homology with field coefficients gives rise to topological arbiters on even dimensional projective spaces, and similarly there are local homological arbiters on D^{2n} for all n , see lemma 1.2 and the discussion following definition 2.1. In this section we use stable homotopy theory to show that there exist local arbiters different from the homological ones in even dimensions ≥ 6 .

Recall that the construction of an uncountable collection of local arbiters on D^4 in section 2 was based on obstructions related to the failure of the Whitney trick in dimension 4. More specifically, this is reflected in the fact that the middle-dimensional (index 2) handles of certain submanifolds of D^4 cannot be embedded disjointly. This argument does not have an analogue in higher dimensions, since the Whitney trick is valid then. Our construction in dimensions 6 and higher, discussed below, is quite different. It is homotopy-theoretic in nature, and the non-embedding results involve handles above the middle dimension. We point out that conversely, this higher-dimensional approach does not work in dimension 4 since the higher homotopy groups of the circle (the attaching region of the submanifolds in D^4) are trivial.

Before outlining a general construction of families of non-homological arbiters on D^{2n} for all $n \geq 4$, we discuss a specific example in dimension 8. (The construction in D^6 requires an additional step, discussed in section 3.3 below.) Consider codimension zero submanifolds $M \subset D^8$ such that $\partial M \cap \partial D^8$ is a regular neighborhood of a standard $S^3 \subset S^7 = \partial D^8$. We will call this $S^3 \subset \partial M$ the attaching 3-sphere of M , and consider *proper* embeddings $M \hookrightarrow D^8$ taking the attaching 3-sphere to ∂D^8 . Our construction of a non-homological arbiter on D^8 relies on the following result.

Lemma 3.1. *There exist submanifolds $(M, S^3) \subset (D^8, S^7)$ such that*

- (1) *for any coefficient ring R the attaching 3-sphere of M is non-trivial in $H_3(M; R)$, and*

(2) *The components of the Hopf link $S^3 \cup S^3 \subset S^7$ do not bound disjoint embeddings of two copies of M into D^8 .*

Proof. Consider a standard 3-sphere $X = S^3 \subset \partial D^8$, let $C = S^3 \times D^4 \times I \subset D^8$ be a collar on its regular neighborhood, and denote by Y a copy pushed into D^8 , $Y = S^3 \times \{0\} \times \{1\}$. Let $\Sigma h: S^4 \rightarrow Y$ be the suspension of the Hopf map $h: S^3 \rightarrow S^2$. Consider a general position map $f: D^5 \rightarrow D^8$ whose restriction $f|_{\partial D^5}$ is Σh . Define M to be the collar C union a regular neighborhood of the image of f in D^8 .

To check (1), first note that the attaching 3-sphere is non-trivial in homology of the abstract CW-complex $Z := S^3 \cup_{\Sigma h} D^5$. The singular set S of a general position map f is 2-dimensional. It follows that the attaching 3-sphere X is non-trivial in $H_3(M)$, indeed since the third homology of the singular set is trivial it follows from the Mayer-Vietoris sequence that $X \neq 0 \in H_3(Z \cup_S \text{cone}(S))$.

Suppose the components S_i^3 of the Hopf link bound disjoint copies $M_i, i = 1, 2$ of M in D^8 . The attaching 4-sphere of the 5-cell in the definition of M is mapped by Σh to the copy Y of S^3 pushed into D^8 . Using the collar structure, it also may be viewed as being mapped by Σh to the actual attaching sphere $X \subset \partial D^8$. Consider the resulting maps $S_i^4 \rightarrow X$.

It is convenient to consider $S^7 = \partial D^8$ as $\mathbb{R}^7$ union the point at infinity, bounding the half space $\mathbb{R}_+^8$. Let A denote the “arrow” map

$$A: M_1 \times M_2 \rightarrow S^7, \quad A(x, y) = \frac{x - y}{|x - y|}.$$

Also consider the composition $\phi: S_1^4 \times S_2^4 \rightarrow S_1^3 \times S_2^3 \rightarrow S^6$ of $\Sigma h \times \Sigma h$ with the arrow map in $\mathbb{R}^7$.

The 5-cells D_i^5 attached to S_i^4 map to M_i , and composing with A one has a map $\psi: D_1^5 \times D_2^5 \rightarrow S^7$. Pick a point $z \in S^6$ which is a regular value both of ϕ and ψ . The preimage of z in $S_1^3 \times S_2^3$ is a single point, and since the fiber of the Hopf map (and of its suspension) is a circle with the Lie group framing, $\phi^{-1}(z)$ is an Arf invariant 1 torus T , the generator of $\Omega_2^{fr} \cong \pi_2^S$, cf [14]. (Related stable homotopy invariants of higher dimensional links have been considered by several authors in the context of link homotopy, see for example [11], [7].) However $\phi^{-1}(z)$ is the boundary of the 3-manifold $M^3 = \psi^{-1}(z) \subset D_1^5 \times D_2^5$. Note that T is codimension 6 in $S^4 \times S^4$, while M^3 is codimension 7 in $D_1^5 \times D_2^5$. Consider the pullback ν under ψ of the normal line bundle to S^6 in S^7 . The 5-cells D_i^5 may be assumed to be mapped near their boundary into the collar $C = S^3 \times D^4 \times I \subset D^8$ as a product $\Sigma h \times Id_I$. Using this product structure, one checks that there is a well-defined limit of ν at the boundary $\partial M = T$, which amounts to a stabilization of the 6-framing of T . This exhibits the Arf invariant 1 torus T as the framed boundary of M , a contradiction. $\square$

Lemma 3.1 easily yields a non-homological arbiter $\mathcal{A}$ on D^8 . This step is analogous to the corresponding part of the proof of theorem 1.3 in dimension 4: consider the decomposition $D^8 = M \cup N$, where M is the manifold constructed in the proof of lemma 3.1, and N is its complement. Given a submanifold $P \subset D^8$ such that M is isotopic to a submanifold of P , set $\mathcal{A}(P) = 1$. On the other hand, if P is isotopic to a submanifold of N , set $\mathcal{A}(P) = 0$. The condition (2) in lemma 3.1 ensures that this consistently defines a partial arbiter on the “cone” (with respect to the partial ordering induced by inclusion) over M and the corresponding cone under N , see figure 4. Now pick any arbiter on D^8 , for example the rational homological arbiter $\mathcal{A}_{\mathbb{Q}}$, and use it to extend $\mathcal{A}$ to all other submanifolds of D^8 . The condition (1) in lemma 3.1 shows that $\mathcal{A}$ is different from the homological arbiters on D^8 .

Suspending the Hopf map further, one gets the analogue of lemma 3.1 and a non-homological arbiter on D^{2n} for any $n > 3$. In fact, the relevant structure used in the proof of the lemma is a non-trivial square in the stable homotopy ring of spheres. As another illustration of this construction, consider the Hopf fibration $H: S^7 \rightarrow S^4$. Following the proof of lemma 3.1 with Σh replaced by $\Sigma^3 H$ (and suspending $\Sigma^3 H$ further), one gets arbiters on D^{2n} , $n \geq 8$. Here at least three suspensions are needed to ensure that the dimension of the singular set S is smaller than the dimension of the attaching sphere of M , so that 3.1 (1) holds. The following theorem summarizes the results discussed above, the case $n = 3$ is proved in section 3.3 below.

Theorem 3.2. *To each non-trivial square in the stable homotopy ring of spheres there is associated a local topological arbiter not induced by homology on D^{2n} for sufficiently large n . In particular, there exist non-homological local arbiters on D^{2n} for each $n \geq 3$.*

3.3. Arbiters on D^6 . Constructing non-homological arbiters on D^6 requires a bit more care. Indeed, note that a direct analogue of the proof of lemma 3.1 (with the actual Hopf map, rather than its suspension) does not hold. (A conceptual explanation for this is provided by Lemma 3.4 below.) Specifically, to recall the notation consider a standard 2-sphere $X = S^2 \subset \partial D^6$, let $C = S^2 \times D^3 \times I \subset D^6$ be a collar on its regular neighborhood, and denote by Y a copy of the 2-sphere pushed into D^6 , $Y = S^2 \times \{0\} \times \{1\}$. Let $h: S^3 \rightarrow Y$ be the Hopf map. Consider a general position map $f: D^4 \rightarrow D^6$ whose restriction $f|_{\partial D^4}$ is h , and define M to be the collar C union a regular neighborhood of the image of f in D^6 .

Analogously to the proof of 3.1, the attaching 2-sphere is non-trivial in homology of $Z := S^2 \cup_h D^4 (= \mathbb{C}P^2)$, and the singular set S of a general position map f is still 2-dimensional. However, in contrast with 3.1, we show in the lemma below that the attaching 2-sphere is in fact trivial in $H_2(M; \mathbb{Z}/2)$ for any map f . To illustrate this point, we consider a specific map $f: (\mathbb{C}P^2, \mathbb{C}P^1) \rightarrow (D^4, S^3)$. Take a Chern class one normal circle bundle B over Y in D^6 . (To be more specific, B is considered as

a subset of the boundary $Y \times S^3$ of a regular neighborhood of Y , and the fiber of $B \rightarrow Y$ over $y \in Y$ is the fiber $h^{-1}(y) \subset \{y\} \times S^3$ of the Hopf fibration h .)

Cone B to the origin of D^6 and also consider the mapping cylinder of the fibration $B \rightarrow Y$, that is the circle fiber over each point $y \in Y$ is coned off to y . (This mapping cylinder gives a map of the Chern class 1 disk bundle into D^6 , bounded by B .) This construction yields a map of CP^2 into D^6 which meets ∂D^6 in the 2-sphere X and the singularities may be arranged to be contained in the suspension Σ of a single fiber F of the circle bundle B . The suspension is formed by the disk fiber bounded by F union with (F coned off to the center of D^6 .) Note that this suspension Σ represents the same class in $H_2(CP^2)$ as $[CP^1]$. Moreover, when Σ is mapped into D^6 it is flattened into a plane, thereby killing the attaching 2-sphere X in $H_2(M; \mathbb{Z})$.

The reader may find it useful to consider a construction analogous to the above for $(\mathbb{R}P^2, \mathbb{R}P^1)$ mapped into (D^3, S^2) where the Chern class 1 bundle B is replaced by the Möbius band. The singularity in this case is formed by a circle, generating $\pi_1(\mathbb{R}P^2)$, flattened in a line through the origin in D^3 .

The following lemma shows that the behavior exhibited in the example above holds (at least with $\mathbb{Z}/2$ coefficients) for *any* general position map f .

Lemma 3.4. *Let $f: CP^2 \rightarrow D^4$ be a general position map taking CP^1 to a 2-sphere in ∂D^4 . Then the class $[CP^1]$ is trivial in $H_2(f(CP^2); \mathbb{Z}/2)$.*

Before proving this lemma, observe that as a consequence the construction given in lemma 3.1 of an arbiter on D^8 fails to give a *non-homological* arbiter on D^6 : since the attaching 2-sphere is trivial in $H_2(M; \mathbb{Z}/2)$, there is a homological obstruction to embedding two copies of M into D^6 forming the Hopf link in ∂D^6 . A non-homological arbiter on D^6 is constructed further below in section 3.6.

Proof of lemma 3.4. Let M be a regular neighborhood of $f(CP^2)$ in D^6 . Attaching another 6-cell to D^6 , consider M as a submanifold of S^6 . Suppose the statement of the lemma is false for this map f , then in particular $[CP^1] \neq 0 \in H_2(\partial M; \mathbb{Z}/2)$. ∂M is a 5-manifold, and there exists [15] a 3-manifold $N^3 \subset \partial M$ representing the Poincaré dual $PD([CP^1]) \in H_3(\partial M; \mathbb{Z}/2)$. Push N outward to get a copy N' which links the 2-sphere Y in S^6 . Consider the arrow map $A: S^2 \times N \rightarrow S^5$, and composing with the Hopf map one has $A': S^3 \times N \rightarrow S^5$. Since the linking number between $Y = S^2$ and N' is 1 (mod 2), the preimage $A'^{-1}(y)$ is a collection of an odd number of circles, each one with the Lie group framing. We will give an argument assuming the preimage $A'^{-1}(y)$ is a single such circle C , the general case is proved analogously.

Since N is disjoint from M , A' extends to a map $D^4 \times N \rightarrow S^5$, and therefore $A'^{-1}(y)$ is null-cobordant: there exists a framed (perhaps non-orientable) surface

$F \subset D^4 \times N^3$, $\partial F = C$. Consider D^4 as the unit ball in $\mathbb{R}^4$; C is a circle with non-trivial framing in $\partial D^4 \times N$. Close off F to get $\overline{F} \subset \mathbb{R}^4 \times N^3$ whose normal bundle ν is trivialized over F but $w_2(\nu) = 1$. The proof of the lemma concludes with the following proposition.

Proposition 3.5. *Suppose $\overline{F}$ is a surface embedded in $N^3 \times \mathbb{R}^4$, where N is a (perhaps non-orientable) 3-manifold. Denoting by ν the normal bundle over $\overline{F}$, the following relation holds between the Stiefel-Whitney classes of ν : if $w_1(\nu) = 0$ then $w_2(\nu) = 0$.*

Proof. Consider the composition g of the embedding e with the projection onto N , $g: \Sigma \hookrightarrow N \times \mathbb{R}^4 \rightarrow N$, and homotope e so that g is an immersion. Denoting by ν' the normal bundle of the immersion g , note that since $\overline{F}$ is codimension one in N , $w_1(\nu') = 0$ implies $w_2(\nu') = 0$. By naturality the same conclusion holds for ν as well. $\square$

3.6. Non-homological arbiters on D^6 . In this section we present a modification of the argument in lemma 3.1 which works in dimension 6. That is, we prove the existence of submanifolds $(M, S^2) \subset (D^6, S^5)$ such that

- (1) for any coefficient ring R the attaching 2-sphere of M is non-trivial in $H_2(M; R)$, and
- (2) The components of the Hopf link $S^2 \cup S^2 \subset S^5$ do not bound disjoint embeddings of two copies of M into D^6 .

The construction of M relies on the map $f: D^4 \rightarrow D^6$ introduced at the beginning of section 3.3. Recall that the singular set $S(f)$ of that map is a 2-sphere in D^4 , representing the same class as $[\mathbb{C}P^1]$ (and the same as the class of the attaching 2-sphere $[X]$) in the second homology of $\mathbb{C}P^2 = S^2 \cup_h D^4$. We will show that a suitable M may be constructed as a regular neighborhood of a map g : punctured $S^2 \times S^2 \rightarrow D^6$. Consider a circle linking the 2-sphere $S(f)$ in D^4 and perform a 1-surgery on D^4 along this circle, the result $D^4 \setminus (S^1 \times D^3) \cup (D^2 \times S^2)$ is a punctured $S^2 \times S^2$. One checks that the 5-dimensional 2-handle $D^2 \times D^3$ (the trace of the surgery) may be embedded in the complement of $f(D^4)$. Denoting by g the resulting map: punctured $S^2 \times S^2 \rightarrow D^6$, note that the singular set of g is the same 2-sphere, $S(g) = S(f)$.

Let M be a regular neighborhood of the image of g , and denote by a, b the generators of $H_2(S^2 \times S^2)$, corresponding respectively to the core and the co-core of the 2-handle. The following relation holds in $H_2(M)$: $[X] = [S(g)] + b$, where X is the attaching sphere of M . It follows that $[X] \neq 0 \in H_2(M)$. As in the beginning of section 3.3, it may be helpful to follow this construction in the half-dimensional real context, where the analogue of g maps a punctured torus into D^3 .

The final observation is that the proof of (2) in lemma 3.1 still goes through when the 5-cells D_i^5 are replaced by an arbitrary orientable punctured 5-manifold attached to the 3-sphere by the suspension of the Hopf map. The same argument in dimension 6 establishes for M the property (2) above. A non-homological arbiter $\mathcal{A}$ is then defined by setting $\mathcal{A}(N) = 1$ for any $N \supset M$, $\mathcal{A}(N) = 0$ for $N \subset D^6 \setminus M$, and extending it to an arbiter on all submanifolds of D^6 .

APPENDIX A. ARBITERS AND SURGERY

An additional property relating the notion of a topological arbiter to 4-dimensional surgery (via the (A, B) -slice reformulation [3, 4]) is stated as axiom (4) below. As in definition 2.1 of a local arbiter, $\mathcal{M}$ denotes $\mathcal{M}_{D^4} = \{(M, \gamma) \mid M \text{ is a codimension zero, smooth, compact submanifold of } D^4, \gamma \subset \partial M, \text{ and } M \cap \partial D^4 \text{ is a tubular neighborhood of an unknotted circle } \gamma \subset S^3\}$. Given $(M', \gamma'), (M'', \gamma'') \in \mathcal{M}$, consider their “double” $D(M', M'') = (S^1 \times D^2 \times I) \cup (M' \cup M'')$ where M', M'' are attached to $S^1 \times D^2 \times \{1\}$ so that γ', γ'' form the Bing double of the core of the solid torus, figure 9. Let γ denote the core of $S^1 \times D^2 \times \{0\}$.

Note that since the components of the Bing double of the attaching curve of a 2-handle $H = D^2 \times D^2$ bound disjoint 0-framed disks in H , there is an embedding $(D(M', M''), \gamma) \subset (D^4, S^3)$ restricting to the given inclusions $(M', \gamma'), (M'', \gamma'') \subset (D^4, S^3)$. In the axiom (4) below, $(D(M', M''), \gamma) \in \mathcal{M}$ denotes *any* embedding $(D(M', M''), \gamma) \subset (D^4, S^3)$ restricting to the given inclusions $M', M'' \subset D^4$. Consider the following condition on a local arbiter:

- (4) Let $(M', \gamma'), (M'', \gamma'') \in \mathcal{M}$ be such that $\mathcal{A}(M', \gamma') = \mathcal{A}(M'', \gamma'') = 1$. Then $\mathcal{A}(D(M', M''), \gamma) = 1$.




FIGURE 9. Bing Double $D(M', M'')$.

Lemma A.1. *If a local arbiter $\mathcal{A}: \mathcal{M} \rightarrow \{0, 1\}$ (definition 2.1) satisfies the axiom (4) above, it provides an obstruction to topological 4-dimensional surgery for the non-abelian free groups.*

Proof. We assume the reader is familiar with the $A - B$ slice formulation [3, 5] of the surgery conjecture. Let $\mathcal{A}$ be an arbiter satisfying (4). Suppose the Borromean Rings $Bor = (l_1, l_2, l_3)$ is an $A - B$ slice link. Consider the given decompositions $D^4 = A_i \cup B_i$, $i = 1, 2, 3$, where the attaching curves α_i, β_i form the Hopf link in S^3 . Renaming the two parts of each decomposition if necessary, using the duality axiom (3) in definition 2.1, one may assume that $\mathcal{A}(A_i, \alpha_i) = 1$ for each i . Since Bor is assumed to be $A - B$ slice, there are disjoint embeddings $f_i: A_i \rightarrow D^4$ such that $f_i(\alpha_i) = l_i$, $i = 1, 2, 3$. Consider the Hopf link $H = (a, b)$, and consider the first two components l_1, l_2 of Bor as the Bing double of the core of a tubular neighborhood of a , and $l_3 = b$. Viewed from this perspective, the submanifolds $f_1(A_1), f_2(A_2)$ combine to give an embedding of a double $f: (D(A_1, A_2), \gamma) \rightarrow (D^4, a)$. By axiom (4), $\mathcal{A}(D(A_1, A_2)) = 1$. Consider the decomposition $D^4 = A \cup B$, where $A = f(D(A_1, A_2))$, $B = D^4 \setminus f(D(A_1, A_2))$. The attaching curves of A, B form the Hopf link a, b . Since $f_3(A_3) \subset B$, by axiom (2) $\mathcal{A}(B) = 1$. Therefore $\mathcal{A}(A), \mathcal{A}(B)$ both equal 1, a contradiction with the axiom (3). $\square$

The $A - B$ -slice problem, a reformulation of the topological 4-dimensional surgery conjecture, was introduced by the first author in [3, 4] and further investigated in [5]. It is believed that the Borromean Rings are not $A - B$ slice, and this would imply that surgery fails for the non-abelian free groups. A stronger conjecture posed in these references asserted that the Borromean Rings are not even a *weakly* $A - B$ slice link. “Weakly $A - B$ slice” means that the embedding data of the submanifolds A_i, B_i is not taken into account. The second author showed in [8] that the Borromean Rings are in fact weakly $A - B$ slice (section 2.9 above has a more detailed discussion of this construction.) This result puts the focus on the actual (rather than “weak”) $A - B$ slice condition, where one keeps track of the embedding of the submanifolds into the 4-ball. The notion of a topological arbiter, additionally satisfying the condition (4) above, axiomatizes the properties which an obstruction to surgery should satisfy in this context. The techniques used in the construction of topological arbiters in section 2 do not give one a sufficient control over their behavior under Bing doubling. Therefore an important open question, relevant for the surgery conjecture, is: *Does there exist an arbiter $\mathcal{A}: \mathcal{M} \rightarrow \{0, 1\}$ satisfying axioms 1 – 4?*

APPENDIX B. MULTI-ARBITERS.

This section introduces two multi-valued generalizations of topological arbiters, and investigates their properties and applications. In particular, a non-homological multi-arbiter is constructed in section B.3, and an application to percolation theory is given in B.4.

Consider a d dimensional manifold M divided into k pieces M_i , $M = \cup_i M_i$, with disjoint interiors, colored $1, 2, \dots, k$. To avoid pathology for PL or smooth manifolds,

we require that each piece is a union of generic cells (dual to a triangulation). Let D_k be such a decomposition of M .

Definition B.1. (“Power set multiarbiters”) A k -arbiter on a manifold M is a function from the set of k -decompositions of M to 2^{2^k} , $f: \{D_k\} \rightarrow 2^{2^k}$, obeying the axioms below. For a subset $S \subset \{1, \dots, k\}$, i.e. $S \in 2^k$, let $|S| \subset M$ be $|S| = \cup_{i \in S} M_i$. Thus $| \cdot |$ depends on D_k (indicated by a superscript below) and f will be interpreted as a rule for assigning 0 or 1 to the submanifolds $|S|$. Let $\overline{S}$ denote the complement of S . The following axioms are imposed on all proper subsets of $\{1, \dots, k\}$, other than the empty set.

- (1) $f(|S|) + f(|\overline{S}|) = 1$.
- (2) $f(|S_1|) = f(|S_2|) = 1 \Rightarrow f(|S_1 \cap S_2|) = 1$.
- (3) If $S \subseteq T$ and $f(|T|) = 1$ then $f(|S|) + f(|T \setminus S|) \geq 1$.
- (4) If D_k^1 and D_k^2 are two k -decompositions so that for $i \in S$ $M_i^1 \subseteq M_i^2$ then $f(|S|^1) = 1 \Rightarrow f(|S|^2) = 1$.
- (5) If there is an isotopy of M carrying D_k^1 to D_k^2 then $f(D_k^1) = f(D_k^2)$.

The intuition is that $f(|S|) = 1$ corresponds to $|S|$ carrying some predetermined homological cycle. The definition can be applied either to closed manifolds M or manifolds with boundary. In the latter case it seems most interesting when a k -decomposition of ∂M is fixed in advance.

To absorb the definition consider a few examples. First let M be the real projective space $\mathbb{R}P^d$, set $k = d$, and define $f(|S|) = 1$ iff $|S|$ carries the non-trivial $\mathbb{Z}/2$ -homology class of $\mathbb{R}P^d$ in dimension cardinality(S) =: $c(S)$. Notice when $d = 2$, one may identify this 2-arbiter with the unique arbiter on $\mathbb{R}P^2$ described in the introduction. Similarly, picking a field F yields d -arbiters on the $2d$ - and $4d$ -dimensional manifolds $\mathbb{C}P^d$ and $\mathbb{H}P^d$.

B.2. A relative example: take a d -cube $[-1, 1]^d$ with a d -decomposition of the boundary with the i th piece equal to the opposite $(d - 1)$ -cubes along the i th coordinate. Set $f(|S|) = 1$ iff

$$H_c(|S|, |S| \cap \partial I^d; F) \longrightarrow H_c(I^d, |S| \cap \partial I^d; F)$$

is onto, where $c = c(S)$ and F is some chosen field. Axioms 1, 2 and 3 are elementary applications of duality. Axiom 4, the “greedy” axiom, is also (literally) natural if one has homological examples in mind. Finally, axiom 5 is topological invariance (the isotopy of axiom 6 is assumed to be identity on boundary in the case of a fixed decomposition on ∂M .) Note that there is a unique homological arbiter (independent of the coefficient field F) on the 3-cube.

Here are a few observations on k -arbiters.

B.3. A non-homological relative multiarbiter on the 3-cube may be constructed as follows. When the cardinality $c(S) = 1$, set $f(|S|) = 0$ if $|S|$ may be isotoped (rel boundary) in I^3 into the collar on $|S| \cap \partial I^3$, and $f(|S|) = 1$ otherwise. Dually, for $c(S) = 2$ set $f(|S|) = 1$ if $|S|$ contains a 2-disk $(D^2, \partial D^2) \subset (|S|, |S| \cap \partial I^3)$ and $f(|S|) = 0$ otherwise. This definition makes it clear that $f(|S|) + f(|\bar{S}|) = 1$ for $c(S) = 1, 2$, implying axiom 1.

Axiom 2 holds since given $S_1 \neq S_2 \subset \{1, 2, 3\}$ of cardinality 2, if both $|S_1|, |S_2|$ contain disks then their intersection is homologically essential (rel boundary) and therefore $|S_1 \cap S_2|$ cannot be isotoped into a collar. To establish (3), suppose $T = \{a, b\}$ and $f(|T|) = 1$. Then $|T|$ contains a disk D as above, which is “colored” by two colors a, b (this is induced by the decomposition $|T| = M_a \cup M_b$ in the notation of definition B.1). This is a relative 2-arbiter on the disk D , and precisely one of the two colors, say a , goes across. This implies $f(|a|) = 1$, concluding the proof of (3).

A similar construction yields non-homological multi-arbiters on higher-dimensional cubes. Start with the homological d -arbiter on I^d with some field coefficients F ($d \geq 3$), discussed in section B.2, and modify it in highest and lowest dimensions: if $c(S) = 1$, set $f(|S|) = 0$ if and only if $|S|$ may be isotoped rel boundary into the collar on $|S| \cap \partial I^d$. If $c(S) = d - 1$ set $f(|S|) = 1$ iff $|S|$ contains a standard $(d - 1)$ -disk $(D^{d-1}, \partial D^{d-1}) \subset (|S|, |S| \cap \partial I^d)$. The axiom check is analogous to that for the 3-cube above.

B.4. Recall the homological k -arbiter mentioned for the k -cube in section B.2: $f(|S|) = 1$ iff

$$H_c(|S|, |S| \cap \partial I^d; \mathbb{Z}_2) \longrightarrow H_c(I^d, |S| \cap \partial I^d; \mathbb{Z}_2)$$

is onto. It is quite easy to see (by induction on dimension) that for at least one singleton $a \in \{1, 2, \dots, k\}$, $f(a) = 1$. A statement in percolation follows immediately. Divide I^d into generic Voronoy cells whose centers are selected by a uniform Poisson process. Color each cell black with probability $= 1/d$. Then the probability that any two opposite faces of I^d are connected by a chain of black cells must be $\geq 1/d$. The proof follows directly from consideration of intersection theory and symmetry.

Consider another version of multiarbiters, whose definition is motivated by Poincaré duality and which is a direct generalization of the $\{0, 1\}$ -valued arbiters (definition 1.1 in the introduction.)

Definition B.5. (“Poincaré multiarbiters”) Let M be a codimension zero submanifold of a d -dimensional manifold W . A *multiarbiter* f associates to M a number $f(M) \in \{0, \dots, d - 1\}$, subject to the axioms:

- (1) If M is ambiently isotopic to M' in W then $f(M) = f(M')$.
- (2) If $M \subset M'$ then $f(M) \leq f(M')$.

- (3) Suppose A, B are codimension zero submanifolds such that $W = A \cup B$, with $A \cap B = \partial A \cap \partial B$. Then $f(A) + f(B) = d - 1$.
- (4) $f(A \cap B) \geq f(A) + f(B) - d$.

In each dimension d , the homological multiarbiter on the projective space $\mathbb{R}P^d$ may be defined by setting $f(M) = \text{maximal } k \text{ such that } H_k(M; \mathbb{Z}/2) \text{ maps onto } H_k(\mathbb{R}P^d; \mathbb{Z}/2)$. The axioms (3)-(4) follow from duality and intersection theory.

The construction in section B.3 may be used to give an example of a non-homological Poincaré multiarbiter on $\mathbb{R}P^3$. Given a codimension zero submanifold $M \subset \mathbb{R}P^3$, set $f(M) = 2$ iff M contains a standard copy of $\mathbb{R}P^2$. Set $f(M) = 0$ iff M can be isotoped into a 3-ball. $f(M)$ is defined to be 1 on all other submanifolds. The duality axiom (3) holds since the complement of $\mathbb{R}P^2$ is a ball, and (4) is a consequence of intersection theory.

Recall from the introduction that if a manifold admits an open book decomposition (for example, any odd-dimensional manifold), then it does not support a topological arbiter. In contrast, any odd-dimensional manifold admits the “trivial” Poincaré multiarbiter f , assigning the value $f(M) = (d - 1)/2$ to each submanifold M . It is easy to see that this is the unique multiarbiter (in the sense of definition B.5) on odd-dimensional spheres. The classification of multiarbiters on other manifolds of dimension ≥ 3 is an interesting open problem.

REFERENCES

- [1] W. Dwyer, *Homology, Massey products and maps between groups*, J. Pure Appl. Algebra 6 (1975), 177-190.
- [2] M. Freedman, *Percolation on the projective plane*, Math. Res. Lett. 4 (1997), 889-894.
- [3] M. Freedman, *A geometric reformulation of four dimensional surgery*, Topology Appl. 24 (1986), 133-141.
- [4] M. Freedman, *Are the Borromean rings (A, B) -slice?*, Topology Appl., 24 (1986), 143-145.
- [5] M. Freedman and X.S. Lin, *On the (A, B) -slice problem*, Topology Vol. 28 (1989), 91-110.
- [6] A. Hatcher, *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [7] U. Koschorke, *Link maps and the geometry of their invariants*, Manuscripta Math. 61 (1988), 383-415.
- [8] V. Krushkal, *A counterexample to the strong version of Freedman’s conjecture*, Ann. of Math. (2) 168 (2008), 675-693.
- [9] V. Krushkal, *Link groups of 4-manifolds*, arXiv:math/0510507.
- [10] T. Lawson, *Open book decompositions for odd dimensional manifolds*, Topology 17 (1978), 189-192.
- [11] W.S. Massey and D. Rolfsen, *Homotopy classification of higher-dimensional links*, Indiana U. Math. J. 34 (1985), 375-391.
- [12] J. Milnor, *Link Groups*, Ann. Math 59 (1954), 177-195.
- [13] F. Quinn, *Open book decompositions, and the bordism of automorphisms*, Topology 18 (1979), 55-73.

- [14] R.E. Stong, Notes On Cobordism Theory, Princeton University Press, 1968.
- [15] R. Thom, *Quelques proprietes globales de varietes differentiables*, Comment. Math. Helv. 28 (1954), 17-86.
- [16] H.E. Winkelnkemper, *Manifolds as open books*, Bull. Amer. Math. Soc. 79 (1973), 45-51.

MICROSOFT STATION Q, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106-6105

E-mail address: `michaelf@microsoft.com`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22904

E-mail address: `krushkal@virginia.edu`